arXiv:1502.06412v4 [stat.ME] 9 Sep 2016

On a non-parametric confidence interval for the regression slope

(Running title: À la Tukey confidence interval for slope)

Róbert Tóth[1] and Ján Somorčík[2*]
[1]Tangent Works, Na Slavíne 1, 81104 Bratislava, Slovakia
[2]Comenius University Bratislava, Mlynská dolina, 84248 Bratislava, Slovakia

Abstract

We investigate an application of the Tukey's methodology in Theil's regression to obtain a confidence interval for the true slope in the straight line regression model with not necessarily normal errors. This specific approach is implemented since 2005 in a package of the software R; however, without any theoretical background. We illustrate by Monte Carlo simulations, that this methodology, unlike the classical Theil's approach based on Kendall's tau, seriously deflates the true confidence level of the resulting interval. We provide also rigorous proofs in case of four data points (in general) and in case of five data points (under some additional conditions); together with a real life methods usage example in the latter case. Summing up, we demonstrate that one should never combine statistical methods without checking the assumptions of their usage and we also give a warning to the already wide community of R users of Theil's regression from various fields of science.

Keywords: Theil's regression; Tukey's confidence interval; Walsh averages; software R.

1 Introduction

The Theil's regression (sometimes referred to as Theil–Sen regression) is a robust non-parametric replacement of the traditional least squares approach to the straight line regression model $Y = \beta_0 + \beta_1 x + \varepsilon$ and also to some more complex linear regression models (the pioneering papers were Theil 1950a,b,c). The Theil's methodology does not require normality of the random errors ε, while being able to provide parameter estimates, tests of linear hypotheses about the parameters, as well as confidence intervals for the parameters (see e.g. Hollander and Wolfe 1999 for a detailed description of the methods).

We focus on the confidence interval (CI) for the true slope β_1. For the software R (R Development Core Team 2010) there exists a package called mblm (Komsta 2013) that includes many tools of Theil's regression. But, surprisingly, when asked for a CI for β_1, the package does not compute the classical Theil's CI for β_1, proposed already in Theil (1950a) and making use of the theory of Kendall's tau. Instead, the package uses a different approach that utilizes without any reference the well-known CI based on the Wilcoxon's signed rank test. In general setting, the CI based on the Wilcoxon's signed rank test has been ascribed to John Tukey (see Hollander and Wolfe 1999 for historical details), who originally developed it to obtain a CI for the true center of symmetry of a symmetric distribution from which we observed a sample of *independent* and identically distributed data. However, it turns out very quickly (see Section 4), that in case of slope estimation in Theil's regression the input data are definitely *not independent*. Therefore, the true confidence level of the resulting interval provided by the package mblm is of question and our paper shows that this negative premonition turns real. We think that it is important to point out and study this issue; and not just for theoretical reasons, i.e. to demonstrate that even a tempting combination of some proven statistical methods may have disastrous consequences if one ignores the assumptions of their usage. Our practical motivation was to alert the community of applied statisticians, because Theil's regression (for its simplicity) and the corresponding R package mblm (for its availability) are rather popular among researchers who apply statistics in other sciences – see e.g. Logan (2011), a textbook for biologists that recommends the package mblm, or this sample of papers reporting the use of the package mblm in microbiology, genetics, chemistry, ecology, forestry, agriculture, hydrology, meteorology and also in behavior analysis: Hunter et al (2012), Carothers et al (2010), Kumari et al (2012), Denys et al (2012), Hunter et al (2013), Lucas et al (2013), Sardans and Peñuelas (2015), Pocewicz et al (2007), Heiskanen et al (2011, 2012), Mueller et al (2014), Arpaci et al (2013), Eastaugh et al (2012), Barroso et al (2015), Cuevas et al (2010), Zottele et al (2010), Puertas Orozco et al (2011), Vannest et al (2013).

Our paper is organized as follows. Section 2 defines the underlying model. In Sections 3 and 4 we provide detailed description of the classical Theil's CI and the CI based on the Tukey's methodology, respectively; with an illustrative application of both methods on a real life dataset of $n = 5$ data points in Section 5. Section 6 shows by means of Monte Carlo simulations that the CI based on the Tukey's

*Corresponding author: somorcik@fmph.uniba.sk

methodology has its true confidence level under the nominal confidence level which is set to the traditional 95% throughout the whole paper. We prove this observation rigorously in case of $n = 4$ data points (Section 8). In Section 7, under some additional conditions, we provide a proof also in the setting of the above-mentioned real life example, i.e. for $n = 5$ data points. For the sake of completeness we treat also the case of $n = 3$ data points in Section 9. Finally, in Section 10 we add some notes on the R package mblm implementation of the CI for the true slope based on the Tukey's methodology. Proofs of the theorems were deferred to the Appendix.

2 The model

For each of the n fixed and distinct points $x_1, x_2, \ldots, x_n$ (values of the predictor x) we observe the value of a random variable Y (response). We get a set of observations $Y_1, Y_2, \ldots, Y_n$, where Y_i is the response at x_i. Without loss of generality we assume that $x_1 < x_2 < \cdots < x_n$. Our linear model has the form

$$Y_i = \beta_0 + \beta_1 x_i + \varepsilon_i, \quad i = 1, 2, \ldots, n,$$

where β_0 (intercept) and β_1 (slope) are unknown parameters. Finally, the unobservable random errors $\varepsilon_1, \varepsilon_2, \ldots, \varepsilon_n$ are iid random variables from a continuous (not necessarily normal) distribution.

3 Theil's confidence interval for slope

The hypothesis

$$H_0 : \beta_1 = \beta^*$$

can be tested using $D_i = Y_i - \beta^* x_i = (\beta_1 - \beta^*)x_i + \beta_0 + \varepsilon_i$. Provided that H_0 is true, the D_i's do not depend on the x_i's, i.e. they do not correlate. Hence, the validity of H_0 can be "measured" e.g. by the sample Kendall's correlation coefficient

$$\tau = \frac{N_c - N_d}{\binom{n}{2}},$$

where N_c is the number of concordant pairs (i.e. pairs of points $[x_i, D_i]$ and $[x_j, D_j]$ such that $(x_i - x_j)(D_i - D_j) > 0$) and N_d is the number of discordant pairs (i.e. pairs of points $[x_i, D_i]$ and $[x_j, D_j]$ such that $(x_i - x_j)(D_i - D_j) < 0$). The test statistic $K = \binom{n}{2}\tau = N_c - N_d$ is known as the Kendall K statistic (see Hollander and Wolfe 1999). By $\mathbb{K}_n$ we denote its distribution under independence of the D_i's from the x_i's. The distribution $\mathbb{K}_n$ has been tabulated (see e.g. Hollander and Wolfe 1999) and implemented in many statistical softwares (see e.g. Wheeler 2009), because it depends just on the sample size n, but not on the distribution of the data. The distribution $\mathbb{K}_n$ is discrete, symmetric and has the support

$$\left\{-\binom{n}{2}, -\binom{n}{2} + 2, -\binom{n}{2} + 4, \ldots, \binom{n}{2} - 4, \binom{n}{2} - 2, \binom{n}{2}\right\},$$

because K has the same parity as $\binom{n}{2}$. A test of the hypothesis H_0 at the significance level α is then

$$\text{reject } H_0 : \beta_1 = \beta^*, \text{ if } |K| \geq k_n(\alpha/2),$$

where $k_n(\alpha/2)$ stands for the upper quantile of the distribution $\mathbb{K}_n$. It should be such an integer that, under H_0, $P(K \geq k_n(\alpha/2)) = \alpha/2$. However, due to the discrete nature of the distribution $\mathbb{K}_n$, an exact equality is virtually impossible. Therefore, we define $k_n(\alpha/2)$ as such a unique integer with the same parity as $\binom{n}{2}$ that

$$P(K \geq k_n(\alpha/2)) \leq \alpha/2 \quad \text{and} \quad P(K \geq k_n(\alpha/2) - 2) > \alpha/2.$$

The consequence is that in general the true probability of the type I error of the above test is bellow the nominal significance level α, because it equals $2 \cdot P(K \geq k_n(\alpha/2))$.

Similar idea leads to a CI for the true slope β_1. Denote

$$S_{ij} = \frac{Y_i - Y_j}{x_i - x_j}, \quad (i < j)$$

the "sample" slope of the line given by the pair of sample points $[x_i, Y_i]$ and $[x_j, Y_j]$. There are $N = \binom{n}{2} = n(n-1)/2$ such slopes. Order them ascendingly and denote the resulting sequence $s_1 < s_2 < \cdots < s_N$ –

since the random ε_i's come from a continuous distribution, we may ignore ties between the sample slopes, because they will happen with zero probability, i.e. we shall assume that there are sharp inequalities between the s_i's. The *Theil's* $1-\alpha$ *confidence interval* for the true slope β_1 is

$$(s_l, s_u), \tag{1}$$

where

$$l = \frac{N - k_n(\alpha/2)}{2} + 1 \quad \text{and} \quad u = \frac{N + k_n(\alpha/2)}{2};$$

see e.g. Hollander and Wolfe (1999). Note that the indices l and u are symmetric in the sense that the same CI is obtained also by taking the l-th slope from bellow a the l-th slope from above, since $l + u = N + 1$. The discrete nature of the distribution $\mathbb{K}_n$ involved causes the true confidence level of the above interval to be typically over $1-\alpha$. The exact value is given by the following theorem.

Theorem 1. *For all k of the form $N-2i$ ($i = 0, 1, \ldots, \lfloor N/4 \rfloor$) put $l = (N-k)/2+1$ and $u = (N+k)/2$. Then the true confidence level of the Theil-type CI (s_l, s_u) is $1 - 2 \cdot P(K \geq k)$ where K is a random variable following $\mathbb{K}_n$.*

Theorem 1 implies that the true confidence level of the Theil's CI (1) equals

$$1 - 2 \cdot P(K \geq k_n(\alpha/2)), \tag{2}$$

which is at least $1-\alpha$.

4 An à la Tukey confidence interval for slope

We start with a brief description of the Tukey's methodology in a general setting (see e.g. Hollander and Wolfe 1999).

Suppose we have some input iid random variables $Z_1, Z_2, \ldots, Z_N$ coming from a continuous symmetric distribution. We compute the so-called Walsh averages $(Z_i + Z_j)/2$ $(i \leq j)$. Now we order the Walsh averages ascendingly (due to the continuity of the underlying distribution, we may ignore ties) and denote the resulting set $w_1 < w_2 < \cdots < w_P$, where $P = \binom{N}{2} + N = N(N+1)/2$. Then the Tukey's $1-\alpha$ CI for the center of symmetry of the true distribution of the Z_i's will be

$$(w_L, w_U), \tag{3}$$

where $U = t_N(\alpha/2)$. The value of L will be "symmetric" in the sense that $L = P - t_N(\alpha/2) + 1$, i.e. one takes the $t_N(\alpha/2)$-th Walsh average from bellow and the $t_N(\alpha/2)$-th from above. Finally, $t_N(\alpha/2)$ denotes the $\alpha/2$ upper quantile of the null distribution of the Wilcoxon's signed rank statistic T^+ having the range $0, 1, 2, \ldots, P$ (see e.g. Hollander and Wolfe 1999 for details). Similarly as with the Theil's CI, the true confidence level of the Tukey's CI is typically strictly above $1-\alpha$.

Now, as the R package mblm does, we apply the Tukey's methodology described above to obtain a CI for the true slope β_1 in Theil's regression. The role of the Z_i's will be played by the set of the slopes of all lines given by all pairs of the data points, i.e. by the set $\{S_{ij}; i < j\}$. We name the resulting interval, i.e. the Tukey's CI based on the slopes S_{ij}, the *à la Tukey confidence interval*. From a particular point of view, it seems to be a good idea to apply the Tukey's approach on the sample slopes S_{ij}, because the Tukey's approach was bred for and is known to perform well in situations of symmetrically distributed data – and it is easy to see that the distributions of our S_{ij}'s are indeed symmetric aroud β_1! Unfortunately, one of the basic assumptions of the Tukey's methodology is independence of the input data, i.e. independence of the Z_i's. However, it is easily seen, that this assumption does not hold for the slopes S_{ij}. Actually, there is functional dependence among the slopes, because, for example, the knowledge of $S_{1,1}, S_{1,2}, \ldots, S_{1,n-1}$ enables us to compute the remaining S_{ij}'s, since

$$S_{ij} = \frac{S_{1j}(x_1 - x_j) - S_{1i}(x_1 - x_i)}{x_i - x_j}.$$

This does not necessarily mean that the à la Tukey CI does not provide at least the nominal confidence level. For example, it may happen that the nominal confidence level is preserved, just the interval is redundantly wide. The best scenario from the à la Tukey CI's point of view is that the interval provides the nominal level of confidence while being narrower than the classical Theil's CI described in Section 3. The real state of affairs will be presented after an illustrative example.

5 A real life example

Let us explain the above-described methods on real data analyzed in Hollander and Wolfe (1999) (Examples 9.1–3) and also in Logan (2011) (Example 8.G), where the following description can be found. Smith (1967) investigated the effects of cloud seeding on rainfall in the Snowy Mountains, Australia. The experiment took place in two areas — the target and the control. Within a year a number of periods were randomly allocated for seeding the target area and additional periods for non-seeding the target area. The total rainfalls $T_{\text{seeded}}, T_{\text{unseeded}}$ in the target and $C_{\text{seeded}}, C_{\text{unseeded}}$ in the control area during the seeding and non-seeding periods were recorded. Within a single year, the impact of seeding was assessed via a double ratio $Y = (T_{\text{seeded}}/C_{\text{seeded}})/(T_{\text{unseeded}}/C_{\text{unseeded}})$ and the experiment was repeated over $n = 5$ years (years denoted by x). The measurements are summarized in Table 1 and depicted on Fig. 1.

Table 1: The cloud seeding experiment — the double ratio Y_i measures the impact of cloud seeding on the rainfall in year x_i

year x_i	1	2	3	4	5
double ratio Y_i	1.26	1.27	1.12	1.16	1.03

Figure 1: The cloud seeding experiment — linear dependence of the double ratio Y_i on time



We adopt the classical straight line regression model $Y_i = \beta_0 + \beta_1 x_i + \epsilon_i$ $(i = 1, 2, \ldots, 5)$. Logan (2011) states that "whilst there may not appear to be any evidence of non-normality…, it could be argued that there are too few observations on which to make meaningful decisions about normality (of the random errors) and it might be safer to not make distributional assumptions". Therefore, the Theil's regression in place of the classical least squares inference is applied. The ordered values of the $N = \binom{5}{2} = 10$ sample slopes $S_{ij} = (Y_i - Y_j)/(x_i - x_j)$ are $s_1 < s_2 < \cdots < s_{10}$: $-.1500$, $-.1300$, $-.0800$, $-.0700$, $-.0575$, $-.0550$, $-.0450$, $-.0333$, $.0100$, $.0400$. The dashed line on Fig. 1 shows the linear trend estimated by the Theil's approach. Its slope is the median of the S_{ij}'s, i.e. $(-0.0575 + (-0.0550))/2 = -0.05625$ and suggests a decreas of the double ratio over time, i.e. a decreas over time of the rainfall increases resulting from the seeding.

Rather than a point estimate, our main concern are the 95% CI's for the true slope. Let us start with the Theil's CI (Section 3). For $\alpha = 5\%$ the appropriate upper quantile $k_5(2.5\%)$ is 10 (see e.g. Table A.30 in Hollander and Wolfe 1999), $l = (10 - 10)/2 + 1 = 1$ and $u = (10 + 10)/2 = 10$. Hence, the resulting CI (1) is (s_1, s_{10}), i.e. it is given by the minimum and the maximum sample slope. Numerically,

$$(s_1, s_{10}) = (-0.15, 0.04)$$

and since the CI contains zero, the negative trend suggested by the point estimate of the true slope does not seem to be significant.

For the the à la Tukey CI, the appropriate upper quantile $t_{10}(2.5\%)$ is 47 (see e.g. Table A.4 in Hollander and Wolfe 1999). The ordered values of the $P = 10 \cdot (10+1)/2 = 55$ Walsh averages $(s_i + s_j)/2$ $(i \le j)$ are $w_1 < w_2 < \cdots < w_{55}$: $-.150, -.140, \ldots, .040$. Further, $U = 47$ and $L = 55 - 47 + 1 = 9$. Therefore, the resulting CI (3) is (w_9, w_{47}), i.e. it is given by the 9-th Walsh average from bellow and the 9-th Walsh average from above – which is the 47-th from bellow, since there are together 55 Walsh

averages. For this dataset,

$$(w_9, w_{47}) = (-0.100, -0.015)$$

and since this CI does not contain zero, it confirms the negative trend suggested by the point estimate of the true slope, i.e. a decrease over time of the rainfall increases resulting from the seeding. Note that, from this point of view, there is a discordance between the Theil's CI and the à la Tukey CI.

The à la Tukey CI $(-0.100, -0.015)$ is reported also in Logan (2011), because the book utilized the R package mblm. For this data, the à la Tukey CI is much narrower then the rather conservative Theil's CI $(-0.15, 0.04)$. This is not just coincidence and in general the à la Tukey CI is not to be trusted because in what follows we show that there is a confidence level issue with it.

6 Monte Carlo study

At the end of Section 4, a positive scenario was hypothesized, that the à la Tukey CI could be narrower than the classical Theil's CI. This was supported also by the real life example about cloud seeding in Section 5. However, this benefit turns out to be worthless because there is a crucial problem with the à la Tukey CI's true confidence level, as can be seen from the results of a Monte Carlo study we have conducted. In Table 2 (except the last column; see below) we provide simulation estimates of the true confidence levels of the à la Tukey CI for the true slope under various settings. The number of data points n changed from 6 to 200. The true values of intercept β_0 and slope β_1 were set to 0 and 1, respectively. The iid random errors ε_i were generated from the normal distribution $N(0, 0.01)$, the Cauchy distribution with location parameter 0 and scale parameter 0.1, or the uniform distribution on the interval $(-0.2, 0.2)$. The motivation for the scale parameters of the distributions was to make the spread of the ε_i's comparable with the spread of the x_i's, i.e. to achieve that the data points $[x_i, Y_i]$ do not produce an ideally straight line, nor resemble a shapeless data cloud. In the part "Evenly spaced x_i's", the x_i's created an equidistant design on the interval $(0, 1)$, more precisely, $x_i = (i-1)/(n-1)$ for $i = 1, 2, \ldots, n$. In the part "Two clusters of evenly spaced x_i's", the design of the experiment consisted of two clusters of evenly spaced points on the subintervals $(0, 1/3)$ and $(2/3, 1)$, more precisely, $x_i = (i-1)/(3(n/2-1))$ for $i = 1, 2, \ldots, n/2$ and $x_i = 2/3 + (i - n/2 - 1)/(3(n/2-1))$ for $i = n/2+1, n/2+2, \ldots, n$. Each figure in Table 2 (except the last column) is based on 10,000 simulations and it is the proportion of times (rounded to three decimal places) the à la Tukey CI covered the true slope β_1. The nominal confidence level was set to 95%.

Just for illustration, the rightmost column of Table 2 contains the true confidence levels of the Theil's CI: these figures are not based on simulations, they have been computed by (2) using the R package SuppDists (Wheeler 2009). Thanks to the distribution-free property of the Theil's CI, these true confidence levels depend just on the number of data n (i.e. not on the design of the x_i's, or on the underlying distribution of the random errors ε_i) and are, of course, always at least as high as 95%.

The main message of Table 2 is that, irrespective of the probability distribution of the random errors, the true confidence level of the à la Tukey CI is strictly below the nominal 95% and decreases rapidly with increasing number of data n in all settings presented in our study. The design of the x_i's does not seem to play an important role either; we tried also some other designs not reported here and the resulting figures were very similar. The Monte Carlo simulations support our suspicion that the method of construction of the à la Tukey CI for the true slope is wrong. In what follows we provide rigorous treatment of the problem in case of $n = 5$, $n = 4$, and $n = 3$ data points.

7 The case of $n = 5$ data points

We are going to examine the true confidence level of the à la Tukey CI for sample size and nominal confidence level as in the real life example about cloud seeding in Section 5, i.e. in case of $n = 5$ data points and nominal confidence level $1 - \alpha = 95\%$. In Section 5 it was derived in detail, that the corresponding Theil's CI (1) and the à la Tukey CI (3) are (s_1, s_{10}) and (w_9, w_{47}), respectively. The following theorem shows their mutual relationship.

Theorem 2. *For $n = 5$, the 95% à la Tukey CI (w_9, w_{47}) is always a subset of the 95% Theil's CI (s_1, s_{10}).*

Theorem 2 itself is not enough to claim that the à la Tukey CI has its true confidence level under 95%. However, by Monte Carlo simulations not reported here we noticed that the à la Tukey CI (w_9, w_{47})

Table 2: True confidence levels of the 95% Theil's CI (computed numerically) and the 95% à la Tukey CI (simulation estimates under various arrangements of the x_i's and various distributions of random errors).

	à la Tukey CI						Theil's CI
	Evenly spaced x_i's			Two clusters of evenly spaced x_i's			
Number of data n	Distribution of random errors: normal	Cauchy	uniform	Distribution of random errors: normal	Cauchy	uniform	
6	.869	.850	.867	.871	.855	.867	.983
10	.804	.773	.793	.804	.777	.800	.953
20	.679	.636	.675	.678	.646	.675	.953
30	.591	.551	.588	.595	.561	.596	.951
40	.533	.494	.540	.530	.499	.541	.952
50	.486	.453	.489	.491	.456	.498	.950
60	.449	.416	.451	.452	.424	.457	.950
70	.425	.388	.427	.426	.396	.429	.951
80	.399	.365	.402	.402	.373	.403	.951
90	.379	.351	.376	.384	.361	.383	.950
100	.357	.340	.364	.369	.343	.365	.950
120	.329	.307	.329	.338	.310	.334	.950
140	.311	.290	.310	.302	.293	.309	.950
160	.294	.267	.295	.303	.278	.296	.950
180	.276	.246	.278	.276	.251	.281	.950
200	.261	.240	.265	.268	.249	.266	.950

happens to be very often the subset of the even narrower Theil-type CI (s_2, s_9). The true confidence level of (s_2, s_9) can be obtained easily: put $l = 2$, $u = 9$, $n = 5$, and $N = 10$, then the notation of Theorem 1 implies that $k = 8$ and the theorem itself gives the true confidence $1 - 2 \cdot P(K \geq 8)$ which can be evaluated e.g. by Table A.30 in Hollander and Wolfe (1999). The approximate result is 91.67%. Therefore, our aim is to show, that the à la Tukey CI (w_9, w_{47}) is "very often" a subset of (s_2, s_9) with the poor confidence level of 91.67%. The consequence will be, that although the true confidence level of the à la Tukey CI (w_9, w_{47}) could be over that of (s_2, s_9), it is definitely under 95%. The next theorem states exact conditions in terms of the s_i's when the above-described desired "very often" inclusion of (w_9, w_{47}) in (s_2, s_9) happens, i.e. conditions when the lower (upper) bound of (w_9, w_{47}) is over (under) the lower (upper) bound of (s_2, s_9).

Theorem 3. *If $n = 5$ then: a) The random event $s_2 \leq w_9$ occurs if and only if $2s_2 \leq s_1 + s_9$. b) The random event $w_{47} \leq s_9$ occurs if and only if $s_2 + s_{10} \leq 2s_9$.*

The following theorem provides an upper bound for the true confidence level $P(\beta_1 \in (w_9, w_{47}))$ of the à la Tukey CI (w_9, w_{47}).

Theorem 4. *Let $n = 5$. Denote*

$$\begin{aligned} p_1 &= P(\beta_1 \in (s_2, s_9) \wedge 2s_2 \leq s_1 + s_9 \wedge s_2 + s_{10} \leq 2s_9) \\ p_2 &= P(\beta_1 \in (s_2, s_{10}) \wedge 2s_2 \leq s_1 + s_9 \wedge 2s_9 < s_2 + s_{10}) \\ p_3 &= P(\beta_1 \in (s_1, s_9) \wedge s_1 + s_9 < 2s_2 \wedge s_2 + s_{10} \leq 2s_9) \\ p_4 &= P(\beta_1 \in (s_1, s_{10}) \wedge s_1 + s_9 < 2s_2 \wedge 2s_9 < s_2 + s_{10}) \end{aligned}$$

Then $P(\beta_1 \in (w_9, w_{47})) \leq p_1 + p_2 + p_3 + p_4$.

Let us discuss Theorem 4 in greater detail. The probability p_1 is simply the true confidence level of (s_2, s_9) together with the probability of the above-discussed "very often" inclusion of (w_9, w_{47}) in (s_2, s_9) with the poor true confidence level 91.67%. This means that p_1 is at most 91.67%. The probabilities p_2, p_3, p_4 are based on the true confidence levels of the slightly wider intervals (s_2, s_{10}), (s_1, s_9), (s_1, s_{10}) and these confidence levels can be high. However, by Theorem 3 it can be seen immediately that the probabilities p_2, p_3, and p_4 reflect only cases when these intervals include also (w_9, w_{47}) whereas (w_9, w_{47}) is not completely included in (s_2, s_9). Monte Carlo simulations suggest that these cases are rare, therefore we hope that p_2, p_3, and p_4 turn out to be low.

All we need to do is to determine the probabilities p_1, p_2, p_3, and p_4. Note that they are given just by the joint probability distribution of the ordered slopes s_i, i.e. one does not need to examine the much more complicated joint probability distribution of the ordered Walsh averages w_i anymore. There are 10! possible orderings of the ten slopes $S_{1,2}, S_{1,3}, \ldots, S_{4,5}$. Denote $B_1, B_2, \ldots, B_{10!}$ the appropriate random events, i.e. each B_i denotes an event that a particular ordering of the slopes happened. Then, p_1 (and similarly p_2, p_3, and p_4) can be decomposed as

$$p_1 = \sum_{i=1}^{10!} P(\{\beta_1 \in (s_2, s_9) \wedge 2s_2 \leq s_1 + s_9 \wedge s_2 + s_{10} \leq 2s_9\} \cap B_i).$$

Let B_i denote, for example, the ordering

$$S_{4,5} < S_{3,5} < S_{2,5} < S_{1,5} < S_{2,3} < S_{2,4} < S_{1,3} < S_{1,4} < S_{3,4} < S_{1,2}. \tag{4}$$

Since

$$S_{ij} = \frac{Y_i - Y_j}{x_i - x_j} = \beta_1 + \frac{\varepsilon_i - \varepsilon_j}{x_i - x_j}, \tag{5}$$

the 9 inequalities in (4) that define the ordering can be rewritten as 9 linear inequalities of the form

$$c_1\varepsilon_1 + c_2\varepsilon_2 + \cdots + c_5\varepsilon_5 < c_0, \tag{6}$$

where $c_0, c_1, c_2, \ldots, c_5$ are constants depending on the x_i's. Further, under B_i the conditions $\beta_1 \in (s_2, s_9)$, $2s_2 \leq s_1 + s_9$ and $2s_9 \geq s_2 + s_{10}$ can be also rewritten as 2+1+1=4 inequalities of the form (6), because under B_i, for example, $s_1 = \beta_1 + \frac{\varepsilon_4 - \varepsilon_5}{x_4 - x_5}$, $s_2 = \beta_1 + \frac{\varepsilon_3 - \varepsilon_5}{x_3 - x_5}$, etc. Therefore,

$$p_1 = \sum_{i=1}^{10!} P(P_i),$$

where $P(P_i)$ is the probability that the random errors vector $(\varepsilon_1, \varepsilon_2, \ldots, \varepsilon_5)^\top$ appears in the 5-dimensional polytope P_i with faces given by the above-mentioned 9+4=13 linear inequalities of the form (6).

Nevertheless, we still have to consider 10! = 3,628,800 polytopes and evaluation of p_2, p_3 and p_4 is going to quadruple this number. Fortunately, most of these polytopes are empty sets because the following theorem implies that a lot of the 10! possible orderings of the slopes are impossible.

Theorem 5. *Let $a < b < c$ be three indices. Then the slope S_{ac} is neither the greatest nor the smallest of the trio S_{ab}, S_{ac}, and S_{bc}.*

An automatized computer inspection of the 10! possible orderings revealed quickly, that only 768 of them conform to Theorem 5, which means that we have to deal just with 768 polytopes to obtain a p_i. At this point we have to set concrete values of the x_i's, because the $P(P_i)$'s depend on them. We decided for the following.

Condition 1. *The x_i's create an equidistant design $x_i = i$ for $i = 1, 2, \ldots, n$.*

As byproducts, this choice of the x_i's has the following pleasant consequences that again reduce the amount of computations.

Theorem 6. *If Condition 1 holds and $n = 5$, then $p_4 = 0$.*

Theorem 7. *If Conditions 1 holds and the distribution of the random errors is symmetric (the latter and fairly common assumption will be posed later), then $p_2 = p_3$.*

Still, it is not easy to evaluate the $P(P_i)$'s under an arbitrary probability distribution of the random errors ε_i. Therefore, we decided for the uniform distribution to make the evaluation easier:

Condition 2. *The probability distribution of the errors ε_i is uniform on the interval $(-1, 1)$.*

Under Condition 2 the probability distribution of the vector $(\varepsilon_1, \varepsilon_2, \ldots, \varepsilon_5)^\top$ is uniform in the 5-dimensional cube with the vertices $[\pm 1, \pm 1, \pm 1, \pm 1, \pm 1]$ and edges of length 2. Therefore, the probabilities $P(P_i)$ reduce to

$$P(P_i) = \frac{V(Q_i)}{2^5},$$

where $V(\cdot)$ denotes volume and the Q_i's are the intersections of the polytopes P_i (each given by a set of above-mentioned 13 linear inequalities) with the cube given by the 10 inequalities

$$\varepsilon_i < 1 \text{ and } \varepsilon_i > -1 \quad (i = 1, 2, \ldots, 5),$$

which means that each Q_i is again a polypote; given by $13 + 10 = 23$ inequalities of the form (6). To evaluate the volumes of such polytopes we used a specialized software Vinci (Büeler and Enge 2003): for each polytope the 23 defining inequalities were passed to Vinci and the computation of the volume was then based on the triangulation of the polytope and computation of determinants.

So we obtained p_1 and p_2 by evaluating the volumes of the above-mentioned $768 + 768$ polytopes and by Theorems 6 and 7 we have

$$p_1 + p_2 + p_3 + p_4 = 0.8107315 + 0.0595787 + 0.0595787 + 0 = 92{,}98889\%.$$

This provides an upper bound for the true confidence level of the interval (w_9, w_{47}), i.e. the true confidence level is under the nominal level 95% (as a Monte Carlo estimate based on 1,000,000 simulations we obtained 87.9%). We note that Condition 1 is just technical, since we are able to evaluate $p_1+p_2+p_3+p_4$ under any particular arrangement of the x_i's. However, Condition 2 about the uniform distribution of the random errors is crucial, because it reduced our computation to evaluation of volumes of polytopes, which could be accomplished by Vinci.

8 The case of $n = 4$ data points

In case of $n = 4$ data points, the 95% à la Tukey CI will be (w_1, w_{21}) which is obviously the same as (s_1, s_6). Its true confidence level can be evaluated by Theorem 1: put $l = 1$, $u = 6$, $n = 4$, $N = 6$, obtain $k = 6$ and the theorem gives the confidence $1 - 2 \cdot P(K \geq 6)$, which can be evaluated e.g. by Table A.30 in Hollander and Wolfe (1999). The approximate result is 91.67% which is definitely under 95%, i.e. the à la Tukey CI does not work correctly in this case either. Note that – unlike the case of $n = 5$ data points – the obtained result 91.67% holds in general, e.g. it is completely independent of the additional Conditions 1 or 2.

The above paragraph also means that in case of $n = 4$ data points the Theil's approach is unable to produce a 95% CI, because the confidence level of (s_1, s_6) (the widest Theil-type interval) is under 95%. From another point of view, the 95% Theil's CI cannot be produced, because $k_4(2.5\%)$ satisfying our definition of the upper quantile value does not exist.

9 The case of $n = 3$ data points

With just $n = 3$ data points at hand, the Theil's approach breaks down, because $k_3(2.5\%)$ does not exist. The same happens to the 95% à la Tukey CI, because the Tukey's methodology does not work for such a low number of data and nominal confidence level of 95% ($t_3(2.5\%)$ does not exist).

10 An R implementation of the à la Tukey confidence interval

As we already noted, the à la Tukey CI for the true slope is implemented in the R package mblm, however, without any reference to a theoretical background. It is available in the CRAN package repository since 2005, but since that time the package documentation has been just noting that the package does not implement the original Theil's CI based on Kendall's tau and it is considered to be implemented in next version of the package. However, it has not been implemented till now (August 2016), despite the fact that already the third version of the package has been released. Nevertheless, the main problem is that the package does not provide any warning about the deflated true confidence level of the intervals produced. The only exceptions are the cases of $n = 4$ and $n = 3$ data points. In case of $n = 4$ data points the package mblm produces a correct warning message, that the requested confidence level is not achievable. However, careful inspection of the package code reveals that it is just a coincidence: in fact, the warning says nothing about the CI for the true *slope*, because it has been invoked by the computation of a CI for the true *intercept* (this CI was not discussed in our paper). For $n = 3$ data points the package mblm produces an error message, however, as before the true reason for the message is a problem with the computation of a CI for the true intercept.

11 Conclusions

We have shown by means of Monte Carlo simulations that the à la Tukey confidence interval for the true slope in the straight line regression model seems to be unable to achieve the nominal confidence level. The loss of interval's confidence does not seem to depend too much on the design of the experiment or on the distribution of the random errors, but becomes very serious with increasing number of data – in all cases with over 160 data points we observed the true confidence level even under 30% instead of the nominal 95%.

In case of $n = 4$ data points we easily obtained also the true confidence level of the à la Tukey confidence interval – the simplicity of the reasoning resulted from the fact that the lower and upper limit of the à la Tukey confidence interval turned out to be some of the original sample slopes. However, in case of $n = 5$ data points the situation was much more complicated: we were able to obtain only an upper bound for the true confidence level and we numerically evaluated this upper bound under the condition of uniformly distributed random errors.

Theoretically, the process of evaluation of the above mentioned upper bound can be adopted to obtain the exact value of the true confidence level of the à la Tukey interval. However, already in case of $n = 5$ data points there are 55 Walsh averages given by the ten slopes S_{ij} and, theoretically, these Walsh averages can be arranged in 55! permutations. These would result in the necessity to evaluate and sum volumes of as much as $55! \approx 1.27 \cdot 10^{73}$ polytopes – a very hard task from the numerical point of view. Similarly as in the evaluation of the p_i's, many of these polytopes could be a priory shown to be of zero volume, but we decided to proceed in a different way: we estimated the true confidence level from above by terms not involving the Walsh averages and showed rather easily that this upper bound is strictly under 95%.

A natural question arises, if the reasoning in case of $n = 5$ data points can be easily adopted or even generalized for larger n. Despite our effort we have not found any positive answer, because the situation complicates dramatically already for $n = 6$.

The à la Tukey confidence interval for the true slope is implemented in the R package mblm without any warning about its deflated true confidence level. The results of our paper show that this functionality of the package (i.e. computation of the confidence interval for the true slope) should not be used, because it tends to provide too liberal interval estimates. We conclude that although the software R is of great help at a great variety of statistical analyses, one has to remember its startup message noting that it "comes with ABSOLUTELY NO WARRANTY".

Apart from the software issue, we provided a simple non-parametric example that an at first glance rather clever combination of some renown statistical methods (Theil's slopes and Tukeys's CI in our case) may yield disastrous results, if one ignores the assumptions of their usage.

Appendix

Proof of Theorem 1

Recall N_c, N_d and the hypothesis H_0 from Section 3. In Theil (1950a) on p. 390, the true confidence level of the Theil-type CI (s_l, s_u) is expressed as

$$1 - 2 \cdot P(N_d \leq l - 1),$$

where the probability is evaluated under H_0 and the result holds even under a more general setting than discussed in our paper. Since $K = N_c - N_d$ and $N_c + N_d = N$, we obtain that $N_d = (N - K)/2$ and

$$1 - 2 \cdot P(N_d \leq l - 1) = 1 - 2 \cdot P\left(\frac{N-K}{2} \leq \left(\frac{N-k}{2} + 1\right) - 1\right) = 1 - 2 \cdot P(K \geq k).$$

□

Proof of Theorem 2

Since the smallest Walsh average w_1 is given by the smallest slope s_1 as $(s_1 + s_1)/2 = s_1$ and the largest Walsh average w_{55} is given by the largest slope s_{10} as $(s_{10} + s_{10})/2 = s_{10}$, we obtain $s_1 = w_1 < w_9$ and $w_{47} < w_{55} = s_{10}$. □

Proof of Theorem 3

Part a): We prove the equivalent statement "$w_9 < s_2$ iff $s_1 + s_9 < 2s_2$". Start with $w_9 < s_2$ and consider the 9 smallest Walsh averages $w_1 < w_2 < \cdots < w_9$. Each of them is of the form $(s_i + s_j)/2$ for

some $i \leq j$ and since $s_2 = (s_2 + s_2)/2$, the assumption $w_9 < s_2$ means that

$$\frac{s_i + s_j}{2} < \frac{s_2 + s_2}{2}. \tag{7}$$

Because $s_1 < s_2 < \cdots < s_{10}$, the sharp inequality (7) immediately implies, that $i = 1$ and the 9 smallest Walsh averages $w_1 < w_2 < \cdots < w_9$ have to be of the form $(s_1 + s_1)/2 < (s_1 + s_2)/2 < \cdots < (s_1 + s_9)/2$. Therefore, the inequality $w_9 < s_2$ can be rewritten as $(s_1 + s_9)/2 < (s_2 + s_2)/2$ and the first part of the proof is complete.

Now, start with $s_1 + s_9 < 2s_2$, i.e. $(s_1 + s_9)/2 < s_2$. Since the 8 Walsh averages $(s_1 + s_1)/2 < (s_1 + s_2)/2 < \cdots < (s_1 + s_8)/2$ are even smaller then $(s_1 + s_9)/2$, we see that there are at least 9 Walsh averages smaller than s_2. Therefore, also the 9-th smallest Walsh averages, i.e. w_9, is smaller than s_2.

Part b): Note that the proof of part a) is based on the natural ordering "the higher slope (or Walsh average), the higher index". Using the reverse ordering "the higher slope (or Walsh average), the lower index" in the proof of part a), one obtains the "symmetric" counterpart of part a), which is part b). □

Proof of Theorem 4

Split the whole probability space into these four disjoint random events:

$$A : s_2 \leq w_9 \wedge w_{47} \leq s_9$$

$$B : s_2 \leq w_9 \wedge s_9 < w_{47}$$

$$C : w_9 < s_2 \wedge w_{47} \leq s_9$$

$$D : w_9 < s_2 \wedge s_9 < w_{47}$$

Denote by U the random event $\{\beta_1 \in (w_9, w_{47})\}$. Note that the minimum and the maximum of all slopes s_i and their Walsh averages w_i are s_1 and s_{10}, respectively. This implies that

$$\begin{aligned}
P(U \cap A) &\leq P(\{\beta_1 \in (s_2, s_9)\} \cap A) = p_1, \\
P(U \cap B) &\leq P(\{\beta_1 \in (s_2, s_{10})\} \cap B) = p_2, \\
P(U \cap C) &\leq P(\{\beta_1 \in (s_1, s_9)\} \cap C) = p_3, \\
P(U \cap D) &\leq P(\{\beta_1 \in (s_1, s_{10})\} \cap D) = p_4,
\end{aligned}$$

where the final equality in each row follows from Theorem 3. Hence, we obtain

$$P(U) = (U \cap A) + P(U \cap B) + P(U \cap C) + P(U \cap D) \leq p_1 + p_2 + p_3 + p_4.$$

□

Proof of Theorem 5

By contradiction, let S_{ac} be the greatest of S_{ab}, S_{ac}, S_{bc} – the case that S_{ac} is the smallest can be treated analogously. By (5) and by noting that $x_a < x_b < x_c$, one observes that the inequality $S_{ac} > S_{ab}$ is equivalent to

$$(\varepsilon_a - \varepsilon_c)(x_a - x_b) > (\varepsilon_a - \varepsilon_b)(x_a - x_c)$$

and $S_{ac} > S_{bc}$ is equivalent to

$$(\varepsilon_a - \varepsilon_c)(x_b - x_c) > (\varepsilon_b - \varepsilon_c)(x_a - x_c).$$

By summing these two inequalities we obtain $(\varepsilon_a - \varepsilon_c)(x_a - x_c) > (\varepsilon_a - \varepsilon_c)(x_a - x_c)$ which is impossible. □

Proof of Theorem 6

Theorem 5 implies that the minimum and maximum sample slopes s_1 and s_{10} are of the form $s_1 = S_{i,i+1}$ and $s_{10} = S_{j,j+1}$ for some distinct i and j from $\{1, 2, 3, 4\}$. Straightforward algebra implies that $S_{i,j} - S_{i+1,j+1} = (s_{10} - s_1)/(i - j)$ under Condition 1, which means that

$$s_{10} - s_1 = |i - j| \cdot |S_{i,j} - S_{i+1,j+1}|. \tag{8}$$

Note that $|i - j| \leq 3$ (because $1 \leq i, j \leq 4$) and if both $S_{i,j}$ and $S_{i+1,j+1}$ belong to $\{s_2, s_3, \ldots, s_9\}$, then one obtains from (8) that

$$s_{10} - s_1 \leq 3(s_9 - s_2). \tag{9}$$

However, summing the inequalities $s_1 + s_9 < 2s_2$ and $2s_9 < s_2 + s_{10}$ appearing in the definition of p_4 yields

$$s_{10} - s_1 > 3(s_9 - s_2),$$

which contradicts (9), i.e. $p_4 = 0$.

It remains to treat the case when not both $S_{i,j}$ and $S_{i+1,j+1}$ belong to $\{s_2, s_3, \ldots, s_9\}$. This happens if and only if $|i - j| = 1$. Without loss of generality, we will suppose that $j = i + 1$, i.e. $s_1 = S_{i,i+1}$, $s_{10} = S_{i+1,i+2}$ and $i \leq 3$.

a) The case when $i \leq 2$. We will show that the inequality

$$2s_9 < s_2 + s_{10} \tag{10}$$

appearing in the definition of p_4 is impossible. Because $s_2 \leq S_{i+2,i+3}$, $S_{i+1,i+3} \leq s_9$ and $s_{10} = S_{i+1,i+2}$, the inequality (10) would imply that $2S_{i+1,i+3} < S_{i+2,i+3} + S_{i+1,i+2}$, which is equivalent to $0 < 0$ under Condition 1.

b) The case when $i = 3$. We will show that the inequality

$$s_1 + s_9 < 2s_2 \tag{11}$$

appearing in the definition of p_4 is impossible. Because $s_2 \leq S_{2,4}$, $S_{2,3} \leq s_9$ and $s_1 = S_{3,4}$, the inequality (11) would imply that $S_{3,4} + S_{2,3} < 2S_{2,4}$, which is equivalent to $0 < 0$ under Condition 1. □

Proof of Theorem 7

Symmetry and independence of the distribution of the ε_i's given by Condition 2, together with the equidistantness of the x_i's given by Condition 1 means that moving from the ε_i's to the "equiprobable" $-\varepsilon_i$'s reverts the ordering of the sample slopes and also the ordering of their Walsh averages, because each sample slope changes symmetrically around β_1 (cf. (5)). It means that, for example, the sample slope with the label s_2 gets the label s_9, or the Walsh average with the label w_9 gets w_{47}, etc. The relationships between the s_i's and w_j's change accordingly: for example, $s_2 \leq w_9$ changes to $w_{47} \leq s_9$. Hence, we observe that the conditions defining p_2 change to conditions defining p_3. □

References

Arpaci A, Eastaugh CS, Vacik H (2013) Selecting the best performing fire weather indices for Austrian ecoregions. Theoretical and Applied Climatology 114:393–406

Barroso LMA, Nascimento M, Nascimento ACC, Fonseca e Silva F, Cruz CD, Bhering LL, de Paula Ferreira R (2015) Metodologia para análise de adaptabilidade e estabilidade por meio de regressão quantílica. Pesquisa Agropecuária Brasileira 50:290–297

Büeler B, Enge A (2003) Vinci. URL http://www.math.u-bordeaux1.fr/∼aenge/index.php?category=software&page=vinci, version 1.0.5. Accessed 17 August 2016

Carothers JM, Goler JA, Kapoor Y, Lara L, Keasling JD (2010) Selecting RNA aptamers for synthetic biology: investigating magnesium dependence and predicting binding affinity. Nucleic Acids Research 38:2736–2747

Cuevas JG, Calvo M, Little C, Pino M, Dassori P (2010) Are diurnal fluctuations in streamflow real? Journal of Hydrology and Hydromechanics 58:149–162

Denys S, Caboche J, Tack K, Rychen G, Wragg J, Cave M, Jondreville C, Feidt C (2012) In vivo validation of the unified barge method to assess the bioaccessibility of arsenic, antimony, cadmium, and lead in soils. Environmental Science & Technology 46:6252–6260

Eastaugh CS, Arpaci A, Vacik H (2012) A cautionary note regarding comparisons of fire danger indices. Natural Hazards and Earth System Science 12:927–934

Heiskanen J, Rautiainen M, Korhonen L, Mõttus M, Stenberg P (2011) Retrieval of boreal forest LAI using a forest reflectance model and empirical regressions. International Journal of Applied Earth Observation and Geoinformation 13:595–606

Heiskanen J, Rautiainen M, Stenberg P, Mõttus M, Vesanto VH, Korhonen L, Majasalmi T (2012) Seasonal variation in MODIS LAI for a boreal forest area in Finland. Remote Sensing of Environment 126:104–115

Hollander M, Wolfe DA (1999) Nonparametric Statistical Methods – 2nd ed. John Wiley & Sons, New York

Hunter WR, Veuger B, Witte U (2012) Macrofauna regulate heterotrophic bacterial carbon and nitrogen incorporation in low-oxygen sediments. The ISME journal 6:2140–2151

Hunter WR, Jamieson A, Huvenne VAI, Witte U (2013) Sediment community responses to marine vs. terrigenous organic matter in a submarine canyon. Biogeosciences 10:67–80

Komsta L (2013) mblm: Median-Based Linear Models. URL http://CRAN.R-project.org/package=mblm, R package version 0.12. Accessed 17 August 2016

Kumari S, Nie J, Chen HS, Ma H, Stewart R, Li X, Lu MZ, Taylor WM, Wei H (2012) Evaluation of gene association methods for coexpression network construction and biological knowledge discovery. PLoS One 7:e50,411

Logan M (2011) Biostatistical Design and Analysis Using R: A Practical Guide. Wiley–Blackwell, Chichester

Lucas RW, Sponseller RA, Laudon H (2013) Controls over base cation concentrations in stream and river waters: A long-term analysis on the role of deposition and climate. Ecosystems 16:707–721

Mueller T, Dressler G, Tucker CJ, Pinzon JE, Leimgruber P, Dubayah RO, Hurtt GC, Böhning-Gaese K, Fagan WF (2014) Human land-use practices lead to global long-term increases in photosynthetic capacity. Remote Sensing 6:5717–5731

Pocewicz A, Vierling LA, Lentile LB, Smith R (2007) View angle effects on relationships between MISR vegetation indices and leaf area index in a recently burned ponderosa pine forest. Remote Sensing of Environment 107:322–333

Puertas Orozco OL, Carvajal Escobar Y, Quintero Angel M (2011) Study of monthly rainfall trends in the upper and middle cauca river basin, Colombia. Dyna–Colombia 78:112–120

R Development Core Team (2010) R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria, URL http://www.R-project.org, ISBN 3-900051-07-0

Sardans J, Peñuelas J (2015) Trees increase their P:N ratio with size. Global ecology and biogeography 24:147–156

Smith EJ (1967) Cloud seeding experiments in Australia. In: Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability, University of California Press, Berkeley, California, USA, vol 5: Weather Modification, pp 161–176

Theil H (1950a) A rank-invariant method of linear and polynomial regression analysis, I. Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen 53:386–392

Theil H (1950b) A rank-invariant method of linear and polynomial regression analysis, II. Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen 53:521–525

Theil H (1950c) A rank-invariant method of linear and polynomial regression analysis, III. Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen 53:1397–1412

Vannest KJ, Davis JL, Parker RI (2013) Single Case Research in Schools: Practical Guidelines for School-Based Professionals. Taylor & Francis, New York

Wheeler B (2009) SuppDists: Supplementary distributions. URL http://CRAN.R-project.org/package=SuppDists, R package version 1.1-8. Accessed 17 August 2016

Zottele F, Toller G, Eccel E (2010) Irri4web: crop water needs definition by webGIS. Italian Journal of Agrometeorology 14:5–14